

Polynomial versus Exponential Growth in Repetition-Free Binary Words

Juhani Karhumäki*

Department of Mathematics and TUCS

University of Turku

20014 Turku

FINLAND

`juhani.karhumaki@cs.utu.fi`

Jeffrey Shallit†

School of Computer Science

University of Waterloo

Waterloo, ON, N2L 3G1

CANADA

`shallit@graceland.uwaterloo.ca`

February 1, 2008

Abstract

It is known that the number of overlap-free binary words of length n grows polynomially, while the number of cubefree binary words grows exponentially. We show that the dividing line between polynomial and exponential growth is $\frac{7}{3}$. More precisely, there are only polynomially many binary words of length n that avoid $\frac{7}{3}$ -powers, but there are exponentially many binary words of length n that avoid $\frac{7}{3}^+$ -powers. This answers an open question of Kobayashi from 1986.

1 Introduction

We are concerned in this paper with problems on combinatorics of words [12, 13].

Let Σ be a finite nonempty set, called an *alphabet*. We consider finite and infinite words over Σ . The set of all finite words over Σ is denoted by Σ^* . The set of all infinite words

*Supported by the Academy of Finland under grant 44087.

†Supported by a grant from NSERC.

(that is, maps from $\mathbb{N}$ to Σ) is denoted by Σ^ω . In this paper we often use a particular class of alphabets, namely

$$\Sigma_k := \{0, 1, \dots, k-1\}.$$

A *morphism* is a map $h : \Sigma^* \rightarrow \Sigma^*$ such that $h(xy) = h(x)h(y)$ for all $x, y \in \Sigma^*$. A morphism may be specified by providing the image words $h(a)$ for all $a \in \Sigma$. If $h : \Sigma^* \rightarrow \Sigma^*$ and $h(a) = ax$ for some letter $a \in \Sigma$, then we say that h is *prolongable* on a , and we can then iterate h infinitely often to get the fixed point $h^\omega(a) := axh(x)h^2(x)h^3(x)\dots$. If there exists an integer k such that the morphism h satisfies $h(a) = k$ for all $a \in \Sigma$, we say it is *k-uniform*. If a morphism is *k-uniform* for some k , then we say it is *uniform*. For more on morphisms, see, for example, [4].

A *square* is a nonempty word of the form xx , as in the English word **murmur**. A *cube* is a nonempty word of the form xxx , as in the Finnish word **kokoko**. An *overlap* is a word of the form $axaxa$, where x is a possibly empty word and a is a single letter, as in the English word **alfalfa**. A word v is a *factor* (sometimes called a *subword*) of a word x if x can be written $x = uvw$ for some words u, w . A word *avoids squares* (resp., cubes, overlaps) if it contains no factor that is a square (resp., cube, overlap). Such words are also called *squarefree* (resp., *cubefree*, *overlap-free*). For example, the English word **square** is squarefree, whereas **squarefree** is not.

It is well-known and easily proved that every word of length 4 or more over a two-letter alphabet contains a square as a factor. However, Thue proved in 1906 [18] that there exist infinite squarefree words over a three-letter alphabet. Thue also proved that the word $\mu^\omega(0) = 0110100110010110\dots$ is overlap-free (and hence cubefree); here μ is the **Thue-Morse morphism** sending $0 \rightarrow 01$ and $1 \rightarrow 10$.

Dejean [5] initiated the study of fractional powers. Let α be a rational number ≥ 1 . Following Brandenburg [2], we say that a nonempty word w is an α -*power* if there exist words $y, y' \in \Sigma^*$ such that $w = y^n y'$, and y' is a prefix of y with $n + |y'|/|y| = \alpha$. For example, the English word **alfalfa** is a $\frac{7}{3}$ -power and the word **ionization** is a $\frac{10}{7}$ -power. If α is a real number, we say that a word w *avoids α -powers* (or is α -*power-free*) if it contains no factor that is a β -power for any rational $\beta \geq \alpha$. We say that a word w *avoids α^+ -powers* (or is α^+ -*power-free*) if it contains no factor that is a β -power for rational $\beta > \alpha$. Thus a word is overlap-free iff it is 2^+ -power-free.

We may enumerate the number of words avoiding various patterns. Brandenburg [2] proved (among other things) that there are exponentially many cubefree binary words; also see Edlin [7]. Restivo and Salemi [15, 16] proved that there exist only polynomially many overlap-free binary words of length n ; in fact they gave an upper bound of $O(n^{\log_2 15})$. The exponent $\log_2 15$ was improved to 1.7 by Kfoury [8] to 1.587 by Kobayashi [10], and to 1.37 by Lepistö [11]. Also see Cassaigne [3].

Overlap-free words avoid 2^+ -powers, and there are only polynomially many over Σ_2 . Cubefree words avoid 3-powers and there are exponentially many over Σ_2 . Kobayashi [9, Problem 6.6] asked the following natural question: at what exponent α (if any) does the number of binary words avoiding α -powers jump from polynomial to exponential? In this paper we prove that the answer is $\frac{7}{3}$. Our proof uses the fact that various structure theorems,

which previously were known for overlap-free words, also hold for any exponent α with $2 < \alpha \leq \frac{7}{3}$.

2 Preliminary lemmas

We begin with some notation and preliminary results. We write $\bar{0} = 1$ and $\bar{1} = 0$. We let μ be the Thue-Morse morphism mentioned in the previous section.

Lemma 1 *Let $t, v \in \Sigma_2^*$. If there exist $c, d \in \Sigma_2$ such that $c\mu(t) = \mu(v)d$, then $d = c$, $t = \bar{c}^n$, and $v = c^n$, where $n = |t| = |v|$.*

Proof. See [1, Lemma 1.7.2]. ■

Lemma 2 *Suppose $t, y \in \Sigma_2^*$ and $\mu(t) = yy$. Then there exists $v \in \Sigma_2^*$ such that $y = \mu(v)$.*

Proof. See [1, Lemma 1.7.3]. ■

Lemma 3 *Let $h : \Sigma^* \rightarrow \Sigma^*$ be a uniform morphism, and let α be a rational number. If w contains an α -power, then $h(w)$ contains an α -power.*

Proof. Suppose w contains an α -power. Then there exist words $s, s' \in \Sigma^+$ and $r, t \in \Sigma^*$ such that $w = rs^n s' t$, where s' is a nonempty prefix of s and $n + |s'|/|s| = \alpha$. Then $h(w) = h(r)h(s)^n h(s')h(t)$. Then $h(w)$ contains the α -power $h(s)^n h(s')$. ■

Note that for arbitrary morphisms the result need not be true (unless α is an integer).

Lemma 4 *Let $w \in \Sigma_2^*$, and suppose $\mu(w)$ contains an α -power. Then w contains a β -power with $\beta \geq \alpha$.*

Proof. Suppose $\mu(w)$ contains an α -power, say $\mu(w) = xy^n y' z$, where $n + |y'|/|y| = \alpha$. There are four cases to consider, based on the parity of $|x|$ and $|y|$.

Case 1: $|x|$ is even and $|y|$ is even. There are two subcases, depending on the parity of $|y'|$.

Case 1a: $|y'|$ is even. Then $|z|$ is even. Then there exist words r, s, s', t , with s' a prefix of s , such that $\mu(r) = x$, $\mu(s) = y$, $\mu(s') = y'$, and $\mu(t) = z$. Then $w = rs^n s' t$, and so w contains the α -power $s^n s'$.

Case 2a: $|y'|$ is odd. Then $|z|$ is odd. Then there exist words r, s, s', t , with s' a prefix of s , and a letter c such that $\mu(r) = x$, $\mu(s) = y$, $\mu(s')c = y'$, and $\bar{c}\mu(t) = z$. Since $|y'|$ is odd, $|y|$ is even, and y' is a prefix of y , it follows that $y'\bar{c}$ is also a prefix of y . Hence $s'c$ is a prefix of s . Then w contains the β -power $s^n s'c$, where

$$\beta = n + \frac{|s'c|}{|s|} = n + \frac{2|s'| + 2}{2|s|} = n + \frac{|y'| + 1}{|y|} > n + \frac{|y'|}{|y|} \geq \alpha.$$

Case 2: $|x|$ is even and $|y|$ is odd. Then there exists a word t such that $\mu(t) = yy$. From Lemma 2 there exists v such that $y = \mu(v)$. But then $|y|$ is even, a contradiction. Thus this case cannot occur.

Case 3: $|x|$ is odd and $|y|$ is even. There are two subcases, depending on the parity of $|y'|$.

Case 3a: $|y'|$ is even. Then $|z|$ is odd. Then there exist words r, s, s', t and letters c, d, e such that $x = \mu(r)c$, $y = \bar{c}\mu(s)d$, $y' = \bar{d}\mu(s')e$, and $z = \bar{e}\mu(t)$. Consideration of the factor yy gives $c = d$. Hence $\mu(w) = \mu(r(cs)^n cs'et)$ and so $w = r(cs)^n cs'et$. Thus w contains the α -power $(cs)^n cs'$, and since s' is a prefix of s , it follows that cs' is a prefix of cs .

Case 3b: $|y'|$ is odd. Then $|z|$ is even. Then we are in the mirror image of case 2a, and the same proof works.

Case 4: $|x|$ is odd and $|y|$ is odd. Then from length considerations we see that there exist words t, v and letters c, d such that $y = c\mu(t) = \mu(v)d$. By Lemma 1, we have $d = c$, $t = \bar{c}^n$, $v = c^n$. Thus $y = c(\bar{c}c)^n$. Since y' is a nonempty prefix of y , we may write $\mu(w) = xy^2ct$ for some word t . Since $|x|$ and $|y|$ are odd, and y ends in c , we must have that cc is the image of letter under μ , a contradiction. Thus this case cannot occur. ■

Theorem 5 *Let $w \in \Sigma_2^*$, and let $\alpha > 2$ be a real number. Then w is α -power-free iff $\mu(w)$ is α -power-free.*

Proof. Combine Lemmas 3 and 4. ■

We remark that Theorem 5 is not true if $\alpha = 2$; for example $w = 01$ contains no square, but $\mu(w) = 0110$ does.

3 A structure theorem for α -power-free words for $2 < \alpha \leq \frac{7}{3}$

Restivo and Salemi [15, 16] proved a beautiful structure theorem for overlap-free binary words. Roughly speaking, it says that any overlap-free word is, up to removal of a short prefix or suffix, the image of another overlap-free word under μ , the Thue-Morse morphism. Perhaps surprisingly, the same sort of structure theorem exists for binary words avoiding α -powers, where α is any real number with $2 < \alpha \leq \frac{7}{3}$.

Theorem 6 *Let x be a word avoiding α -powers, with $2 < \alpha \leq \frac{7}{3}$. Let μ be the Thue-Morse morphism. Then there exist u, v, y with $u, v \in \{\epsilon, 0, 1, 00, 11\}$ and a word $y \in \Sigma_2^*$ avoiding α -powers, such that $x = u\mu(y)v$.*

Proof. We prove the result by induction on $|x|$. If $|x| \leq 2$, then the factorizations can be chosen as shown in the following table.

x	u	y	v
ϵ	ϵ	ϵ	ϵ
0	0	ϵ	ϵ
1	1	ϵ	ϵ
00	00	ϵ	ϵ
01	ϵ	0	ϵ
10	ϵ	1	ϵ
11	11	ϵ	ϵ

Now suppose the claim is true for all x with $|x| < k$. We prove it for $|x| = k$. Let x be α -power-free with $|x| \geq 3$. Write $x = az$ with $a \in \Sigma_2$ and $z \in \Sigma_2^*$. Since x is α -power-free, so is z . Since $|z| < |x|$, by induction there exist $u', v' \in \{\epsilon, 0, 1, 00, 11\}$ and a α -power-free word y' such that $z = u'\mu(y')v'$.

Now there are several cases to consider.

Case 1: $u' = \epsilon$ or $u' = a$. Then we may write $x = u\mu(y)v$ with $(u, y, v) = (au', y', v')$.

Case 2: $u' = \bar{a}$. Then $x = u\mu(y)v$ with $(u, y, v) = (\epsilon, ay', v')$. Since x is α -power-free, so is $\mu(ay')$, and hence, by Theorem 5, so is ay' .

Case 3: $u' = aa$. Then x begins with $aaa = a^3$, and so x does not avoid α -powers.

Case 4: $u' = \bar{a}\bar{a}$. Then $x = a\bar{a}\bar{a}\mu(y')v'$.

Case 4.a: $|y'| = 0$. Then $x = a\bar{a}\bar{a}v'$. If $v' = \epsilon$ (resp., $v' = a$, $v' = aa$), then we can write $x = u\mu(y)v$ with $(u, y, v) = (\epsilon, a, \bar{a})$ (resp., $(u, y, v) = (\epsilon, a\bar{a}, \epsilon)$, $(u, y, v) = (\epsilon, a\bar{a}, a)$). Otherwise, if $v' = \bar{a}$ or $v' = \bar{a}\bar{a}$ then x contains $\bar{a}\bar{a}\bar{a} = \bar{a}^3$, and so x does not avoid α -powers.

Case 4.b: $|y'| \geq 1$. There are two cases to consider.

Case 4.b.i: $y' = ay''$. There are several cases to consider.

Case 4.b.i.1: $|y''| = 0$. Then $y' = a$ and $x = a\bar{a}\bar{a}a\bar{a}v'$.

If $v' = \epsilon$ (resp., $v' = a$, $v' = aa$, $v' = \bar{a}$), then we can write $x = u\mu(y)v$ with $(u, y, v) = (\epsilon, a\bar{a}, \bar{a})$ (resp., $(u, y, v) = (\epsilon, a\bar{a}\bar{a}, \epsilon)$, $(u, y, v) = (\epsilon, a\bar{a}\bar{a}, a)$, $(u, y, v) = (\epsilon, a\bar{a}, \bar{a}\bar{a})$). Otherwise, if $v' = \bar{a}\bar{a}$, then x contains $\bar{a}\bar{a}\bar{a} = \bar{a}^3$, and so x does not avoid α -powers.

Case 4.b.i.2: $|y''| \geq 1$. If $y'' = ay'''$, then $x = a\bar{a}\bar{a}a\bar{a}\bar{a}\mu(y''')v'$, so x contains the $\frac{5}{2}$ -power $\bar{a}a\bar{a}a\bar{a}$. If $y'' = \bar{a}y'''$, then $x = a\bar{a}\bar{a}a\bar{a}\bar{a}\mu(y''')v'$, so x contains the $\frac{7}{3}$ -power $a\bar{a}\bar{a}a\bar{a}\bar{a}$.

Case 4.b.ii: $y' = \bar{a}y''$. Then $x = a\bar{a}\bar{a}\bar{a}a\mu(y'')v'$. Thus x contains $\bar{a}\bar{a}\bar{a} = \bar{a}^3$.

Our proof by induction is now complete.

■

The decomposition in Theorem 6 is actually unique if $|x| \geq 7$. As this requires more tedious case analysis and is not crucial to our discussion, we do not prove this here.

We also note that the role of $\frac{7}{3}$ in Theorem 6 is crucial, since no word $\cdots 0110110 \cdots$ can be factorized in the stated form.

4 Polynomial upper bound on the number of $\frac{7}{3}$ -power-free words

Theorem 6 has the following implication. Let $x = x_0$ be a nonempty binary word that is α -power-free, with $2 < \alpha \leq \frac{7}{3}$. Then by Theorem 6 we can write $x_0 = u_1\mu(x_1)v_1$ with $|u_1|, |v_1| \leq 2$. If $|x_1| \geq 1$, we can repeat the process, writing $x_1 = u_2\mu(x_2)v_2$. Continuing in this fashion, we obtain the decomposition $x_i = u_i\mu(x_i)v_i$ until $|x_{t+1}| = 0$ for some t . Then

$$x_0 = u_1\mu(u_2) \cdots \mu^{t-1}(u_{t-1})\mu^t(x_t)\mu^{t-1}(v_{t-1}) \cdots \mu(v_2)v_1.$$

Then from the inequalities $1 \leq |x_t| \leq 4$ and $2|x_i| \leq |x_{i-1}| \leq 2|x_i| + 4$, $1 \leq i \leq t$, an easy induction gives $2^t \leq |x| \leq 2^{t+3} - 4$. Thus $t \leq \log_2 |x| < t + 3$, and so

$$\log_2 |x| - 3 < t \leq \log_2 |x|. \quad (1)$$

There are at most 5 possibilities for each u_i and v_i , and there are at most 22 possibilities for x_t (since $1 \leq |x_t| \leq 4$ and x_t is α -power-free). Inequality (1) shows there are at most 3 possibilities for t . Letting $n = |x|$, we see there are at most $3 \cdot 22 \cdot 5^{2 \log_2 n} = 66n^{\log_2 25}$ words of length n that avoid α -powers. We have therefore proved

Theorem 7 *Let $2 < \alpha \leq \frac{7}{3}$. There are $O(n^{\log_2 25}) = O(n^{4.644})$ binary words of length n that avoid α -powers.*

We have not tried to optimize the exponent in Theorem 7. Probably it can be made significantly smaller.

5 Exponential lower bound on the number of $\frac{7}{3}^+$ -power-free words

In this section we prove that there are exponentially many binary words of length n avoiding $\frac{7}{3}^+$ -powers.

Define the 21-uniform morphism $h : \Sigma_4^* \rightarrow \Sigma_2^*$ as follows:

$$\begin{aligned} h(0) &= 011010011001001101001 \\ h(1) &= 100101100100110010110 \\ h(2) &= 100101100110110010110 \\ h(3) &= 011010011011001101001. \end{aligned}$$

We first show

Lemma 8 *Let w be any squarefree word over Σ_4 . Then*

- (i) $h(w)$ contains no square yy with $|y| > 13$; and
- (ii) $h(w)$ contains no $\frac{7}{3}^+$ -powers.

Proof. We first prove (i). We argue by contradiction. Let $w = a_1a_2 \cdots a_n$ be a squarefree word such that $h(w)$ contains a square, i.e., $h(w) = xy yz$ for some $x, z \in \Sigma_4^*$, $y \in \Sigma_4^+$. Without loss of generality, assume that w is a shortest such word, so that $0 \leq |x|, |z| < 21$.

Case 1: $|y| \leq 42$. In this case we can take $|w| \leq 5$. To verify that $h(w)$ contains no squares yy with $|y| > 13$, it therefore suffices to check the image of each of the 264 squarefree words in Σ_4^5 .

Case 2: $|y| > 42$. First, we observe the following facts about h .

Fact 9 (i) *Suppose $h(ab) = th(c)u$ for some letters $a, b, c \in \Sigma_4$ and words $t, u \in \Sigma_2^*$. Then this inclusion is trivial (that is, $t = \epsilon$ or $u = \epsilon$) or u is not a prefix of $h(d)$ for any $d \in \Sigma_4$.*

(ii) *Suppose there exist letters $a, b, c \in \Sigma_4$ and words $s, t, u, v \in \Sigma_2^*$ such that $h(a) = st$, $h(b) = uv$, and $h(c) = sv$. Then either $a = c$ or $b = c$.*

Proof.

- (i) This can be verified with a short computation.
- (ii) This can also be verified with a short computation. If $|s| \geq 11$, then no two images of distinct letters share a prefix of length 11. If $|s| \leq 10$, then $|t| \geq 11$, and no two images of distinct letters share a suffix of length 11.

■

Now we resume the proof of Lemma 8. For $i = 1, 2, \dots, n$ define $A_i = h(a_i)$. Then if $h(w) = xy yz$, we can write

$$h(w) = A_1A_2 \cdots A_n = A'_1A''_1A_2 \cdots A_{j-1}A'_jA''_jA_{j+1} \cdots A_{n-1}A'_nA''_n$$

where

$$\begin{aligned} A_1 &= A'_1A''_1 \\ A_j &= A'_jA''_j \\ A_n &= A'_nA''_n \\ x &= A'_1 \\ y &= A''_1A_2 \cdots A_{j-1}A'_j = A''_jA_{j+1} \cdots A_{n-1}A'_n \\ z &= A''_n, \end{aligned}$$

and $|A'_1|, |A''_j| > 0$. See Figure 1.

A'_1	A''_1					A'_j	A''_j					A'_n	A''_n
A_1	A_2	$\cdots$	A_{j-1}	A_j	A_{j+1}	$\cdots$	A_{n-1}	A_n					
x	y				y				z				

Figure 1: The word $xyyz$ within $h(w)$

If $|A'_1| > |A''_j|$, then $A_{j+1} = h(a_{j+1})$ is a factor of $A'_1 A_2$, hence a factor of $A_1 A_2 = h(a_1 a_2)$. Thus we can write $A_{j+2} = A'_{j+2} A''_{j+2}$ with

$$A'_1 A_2 = A'_j A_{j+1} A'_{j+2}.$$

See Figure 2.

$y =$	A_1''	A_2	$\cdots$	A_{j-1}	A_j'	
$y =$	A_j''	A_{j+1}	A_{j+2}'	$\cdots$	A_{n-1}	A_n'

Figure 2: The case $|A'_1| > |A''_j|$

But then, by Fact 9 (i), either $|A''_j| = 0$, or $|A'_{j+2}| = 0$ (so $|A'_1| = |A''_j|$), or A'_{j+2} is not a prefix of any $h(d)$. All three conclusions are impossible.

If $|A'_1| < |A''_j|$, then $A_2 = h(a_2)$ is a factor of $A''_j A_{j+1}$, hence a factor of $A_j A_{j+1} = h(a_j a_{j+1})$. Thus we can write $A_3 = A'_3 A''_3$ with

$$A'_1 A_2 A'_3 = A'_j A_{j+1}.$$

See Figure 3.

$y =$	A'_1	A_2	A'_3	$\cdots$	A_{j-1}	A'_j
$y =$	A''_j	A_{j+1}	$\cdots$	A_{n-1}	A'_n	

Figure 3: The case $|A'_1| < |A''_j|$

By Fact 9 (i), either $|A'_1| = 0$ or $|A'_j| = 0$ (so $|A'_1| = |A''_j|$) or A'_3 is not a prefix of any $h(d)$. Again, all three conclusions are impossible.

Therefore $|A'_1| = |A''_j|$. Hence $A'_1 = A''_j$, $A_2 = A_{j+1}$, $\dots$, $A_{j-1} = A_{n-1}$, and $A'_j = A'_n$. Since h is injective, we have $a_2 = a_{j+1}, \dots, a_{j-1} = a_{n-1}$. It also follows that $|y|$ is divisible by 21 and $A_j = A'_j A''_j = A'_n A'_1$. But by Fact 9 (ii), either (1) $a_j = a_n$ or (2) $a_j = a_1$.

In the first case, $a_2 \cdots a_{j-1} a_j = a_{j+1} \cdots a_{n-1} a_n$, so w contains the square $(a_2 \cdots a_{j-1} a_j)^2$, a contradiction. In the second case, $a_1 \cdots a_{j-1} = a_j a_{j+1} \cdots a_{n-1}$, so w contains the square $(a_1 \cdots a_{j-1})^2$, a contradiction.

This completes the proof of part (i).

It now remains to prove (ii). If $h(w)$ contains a $\frac{7}{3}^+$ -power yyy' , then it contains a square, and by part (i) we know that $|y| \leq 13$. We may assume that $|y'| \leq \lceil \frac{1}{3} \cdot 13 \rceil = 5$, so $|yyy'| \leq 31$. Hence we need only check the image of all 36 squarefree words in Σ_4^3 to ensure they do not contain any $\frac{7}{3}^+$ -power. We leave this computation to the reader.

■

Next, define the substitution $g : \Sigma_3^* \rightarrow 2^{\Sigma_4^*}$ as follows:

$$\begin{aligned} g(0) &= \{0, 3\} \\ g(1) &= \{1\} \\ g(2) &= \{2\}. \end{aligned}$$

Let $|w|_a$ denote the number of occurrences of the letter a in the word w . We prove

Lemma 10 *Let $w \in \Sigma_3^*$ be any squarefree word. Then $h(g(w))$ is a language of 2^r words over Σ_2 , where $r = |w|_0$, and moreover these words are of length $21|w|$ and avoid $\frac{7}{3}^+$ -powers.*

Proof. Let w be a squarefree word over Σ_3 . Then $g(w)$ is a language over Σ_4 , and we claim that each word $x \in g(w)$ is squarefree. For suppose $x \in g(w)$ and x contains a square, say $x = tuuv$ for some words $t, v \in \Sigma_4^*$, and $u \in \Sigma_4^+$. Define the morphism f where $f(0) = f(3) = 0$, $f(1) = 1$, and $f(2) = 2$. Then $f(x) = w$ and $f(x)$ contains the square $f(u)f(u)$, a contradiction. It now follows from Lemma 8 that x avoids $\frac{7}{3}^+$ -powers. ■

Finally, we obtain

Theorem 11 *Let C_n be the number of binary words of length n that are $\frac{7}{3}^+$ -power-free. Then $C_n = \Omega(\gamma^n)$, where $\gamma = 2^{1/63} \doteq 1.011$.*

Proof. Take any squarefree word x of length m over Σ_3 . There must exist a symbol $a \in \Sigma_3$ such that a occurs at least $\lceil m/3 \rceil$ times in x . By replacing each symbol b in x with $b - a \pmod{3}$, we get a squarefree word x' with at least $\lceil m/3 \rceil$ occurrences of 0.

Now consider $h(g(x'))$. We get at least $2^{m/3}$ words of length $21m$, and each word is $\frac{7}{3}^+$ -power-free. Write $n = 21m - k$, where $0 \leq k < 21$. By what precedes, there are at least $2^{n/63}$ words of length $21m$ that are $\frac{7}{3}^+$ -power-free. Thus there are at least $2^{-k} 2^{n/63} \geq 2^{-21} 2^{n/63}$ words of length n with the desired property. ■

We have not tried to optimize the value of γ . It can be improved slightly in several ways: for example, by starting with a squarefree word over Σ_3 with a higher proportion of 0's; see [17].

For an upper bound on C_n , we may reason as follows: if w is a word avoiding $\frac{7}{3}^+$ -powers, then w certainly has no occurrences of either 000 or 111. Let E_n denote the number of binary words of length n avoiding both 000 and 111. Then $C_n \leq E_n$ and it is easy to see that

$$E_n = E_{n-1} + E_{n-2} \quad (2)$$

for $n \geq 3$. Now the characteristic polynomial of the recursion (2) is $x^2 - x - 1$, and the dominant zero of this polynomial is $(1 + \sqrt{5})/2 \doteq 1.62$. By well-known properties of linear recurrences we get $E_n = O(1.62^n)$.

This procedure may be automated. Noonan and Zeilberger [14] have written a Maple package `DAVID_IAN` that allows one to specify a list L of forbidden words, and computes the generating function enumerating words avoiding members of L . We used this package for a list L of 58 words of length ≤ 24 :

$$000, 111, 01010, 10101, \dots, 110110010011011001001101$$

including words of the form $x^{(1+\lceil 7|x|/3 \rceil)/|x|}$ for $1 \leq |x| \leq 10$. (Words for which shorter members of L are factors can be omitted.) We obtained a characteristic polynomial of degree 39 with dominant root $1.22990049\dots$. Therefore we have shown

Theorem 12 *The number C_n of binary words of length n avoiding $\frac{7}{3}^+$ -powers satisfies $C_n = O(1.23^n)$.*

6 Avoiding arbitrarily large squares

Dekking [6] proved that every infinite overlap-free binary word must contain arbitrarily large squares. He also proved that there exists an infinite cube-free binary word that avoids squares xx with $|x| \geq 4$. Furthermore the number 4 is best possible, since every binary word of length ≥ 30 contains a cube or a square xx with $|x| \geq 3$.

This leads to the following natural question: what is the largest exponent α such that every infinite α -power-free binary word contains arbitrarily large squares? From Dekking's results we know $2 < \alpha < 3$. The answer is given in the following theorem.

Theorem 13 (i) *Every infinite $\frac{7}{3}$ -power-free binary word contains arbitrarily large squares.*

(ii) *There exists an infinite $\frac{7}{3}^+$ -power-free binary word such that each square factor xx satisfies $|x| \leq 13$.*

Proof. For (i), let $\mathbf{w}$ be an infinite $\frac{7}{3}$ -power-free binary word. By Theorem 6 and Eq. (1), any prefix of $\mathbf{w}$ of length 2^{n+5} contains $\mu^{n+2}(0)$ as a factor. But $\mu^{n+2}(0) = \mu^n(0110)$, so any prefix of length 2^{n+5} contains the square factor xx with $x = \mu^n(1)$.

For (ii), from Theorem 8 it follows that if $\mathbf{w}$ is an infinite square-free word over Σ_4 , and h is the morphism defined in § 5, then $h(\mathbf{w})$ has the desired properties. ■

We note that the number 13 in Theorem 13 (ii) is not best possible. A forthcoming paper examines this question in more detail.

7 Numerical Results

Let A_n (resp., B_n , C_n , D_n) denote the number of overlap-free words (resp., $\frac{7}{3}$ -power-free words, $\frac{7}{3}^+$ -power-free words, cubefree words) over the alphabet Σ_2 . We give here the values of these sequences for $0 \leq n \leq 28$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
A_n	1	2	4	6	10	14	20	24	30	36	44	48	60	60	62	72	82
B_n	1	2	4	6	10	14	20	24	30	40	48	56	64	76	82	92	106
C_n	1	2	4	6	10	14	20	30	38	50	64	86	108	136	178	222	276
D_n	1	2	4	6	10	16	24	36	56	80	118	174	254	378	554	802	1168

n	17	18	19	20	21	22	23	24	25	26	27	28
A_n	88	96	112	120	120	136	148	164	152	154	148	162
B_n	124	142	152	172	192	210	220	234	256	284	308	314
C_n	330	408	500	618	774	962	1178	1432	1754	2160	2660	3292
D_n	1716	2502	3650	5324	7754	11320	16502	24054	35058	51144	74540	108664

8 Acknowledgments

We would like to thank Ming-wei Wang, who showed the first author the usefulness of Fact 9 (i).

References

- [1] J.-P. Allouche and J. Shallit. *Automatic Sequences: Theory, Applications, Generalizations*. Cambridge University Press, 2003. In press.
- [2] F.-J. Brandenburg. Uniformly growing k -th power-free homomorphisms. *Theoret. Comput. Sci.* **23** (1983), 69–82.
- [3] J. Cassaigne. Counting overlap-free binary words. In P. Enjalbert, A. Finkel, and K. W. Wagner, editors, *STACS 93, Proc. 10th Symp. Theoretical Aspects of Comp. Sci.*, Vol. 665 of *Lecture Notes in Computer Science*, pp. 216–225. Springer-Verlag, 1993.
- [4] C. Choffrut and J. Karhumäki. Combinatorics of words. In G. Rozenberg and A. Salomaa, editors, *Handbook of Formal Languages*, Vol. 1, pp. 329–438. Springer-Verlag, 1997.
- [5] F. Dejean. Sur un théorème de Thue. *J. Combin. Theory. Ser. A* **13** (1972), 90–99.
- [6] F. M. Dekking. On repetitions of blocks in binary sequences. *J. Combin. Theory. Ser. A* **20** (1976), 292–299.

- [7] A. E. Edlin. The number of binary cube-free words of length up to 47 and their numerical analysis. *J. Diff. Equations Appl.* **5** (1999), 353–354.
- [8] A.-J. Kfoury. A linear-time algorithm to decide whether a binary word contains an overlap. *RAIRO Inform. Théor. App.* **22** (1988), 135–145.
- [9] Y. Kobayashi. Repetition-free words. *Theoret. Comput. Sci.* **44** (1986), 175–197.
- [10] Y. Kobayashi. Enumeration of irreducible binary words. *Disc. Appl. Math.* **20** (1988), 221–232.
- [11] A. Lepistö. A characterization of 2^+ -free words over a binary alphabet. Master’s thesis, University of Turku, Finland, 1995.
- [12] M. Lothaire. *Combinatorics on Words*, Vol. 17 of *Encyclopedia of Mathematics and Its Applications*. Addison-Wesley, 1983.
- [13] M. Lothaire. *Algebraic Combinatorics on Words*, Vol. 90 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, 2002.
- [14] J. Noonan and D. Zeilberger. The Goulden-Jackson cluster method: extensions, applications and implementations. *J. Differ. Equations Appl.* **5** (1999), 355–377.
- [15] A. Restivo and S. Salemi. On weakly square free words. *Bull. European Assoc. Theor. Comput. Sci.*, No. 21, (October 1983), 49–56.
- [16] A. Restivo and S. Salemi. Overlap free words on two symbols. In M. Nivat and D. Perrin, editors, *Automata on Infinite Words*, Vol. 192 of *Lecture Notes in Computer Science*, pp. 198–206. Springer-Verlag, 1985.
- [17] Y. Tarannikov. The minimal density of a letter in an infinite ternary square-free word is $0.2746\dots$. *J. Integer Sequences* **5** (2002), 02.2.2 (electronic). <http://www.math.uwaterloo.ca/JIS/VOL5/Tarannikov/tarann7.html>
- [18] A. Thue. Über unendliche Zeichenreihen. *Norske vid. Selsk. Skr. Mat. Nat. Kl.* **7** (1906), 1–22. Reprinted in *Selected Mathematical Papers of Axel Thue*, T. Nagell, editor, Universitetsforlaget, Oslo, 1977, pp. 139–158.